ELSEVIER

European Journal of Pharmacology 279 (1995) 257–258

ejp

Author index

Advenier, C., see Bétoin, F. (279) 241
Akahane, A., see Terai, T. (279) 217
Akkermans, L.M.A., see Briejer, M.R. (279) 123
Amann, R., Schuligoi, R., Lanz, I. and Donnerer, J.
Histamine-induced edema in the rat paw – effect of capsaicin denervation and a CGRP receptor antagonist (279) 227
Ando, T., see Terai, T. (279) 217

Baird, A.W., see Keely, S.J. (279) 203
Barr, G., see Franco-Cereceda, A. (279) 105
Bertha, C.M., see Bowen, W.D. (279) R1
Bétoin, F., Advenier, C., Fardin, V., Wilcox, G., Lavarenne, J. and Eschalier, A.
In vitro and in vivo evidence for a tachykinin NK_1 receptor antagonist effect of vapreotide, an analgesic cyclic analog of somatostatin (279) 241
Bowen, W.D., Vilner, B.J., Williams, W., Bertha, C.M., Kuehne, M.E. and Jacobson, A.E.
Ibogaine and its congeners are σ_2 receptor-selective ligands with moderate affinity (279) R1
Briejer, M.R., Akkermans, L.M.A., Lefebvre, R.A. and Schuurkes, J.A.J.
Novel 5-HT_2-like receptor mediates neurogenic relaxation of the guinea-pig proximal colon (279) 123
Bruning, H.W., see Van Ree, J.M. (279) 187

Chapman, A.G., see Millan, M.H. (279) 75
Chatelain, P., see Hodeige, D. (279) 25

Dhawan, B.N., see Kumar, A. (279) 1
Donnerer, J., see Amann, R. (279) 227

Eschalier, A., see Bétoin, F. (279) 241

Fardin, V., see Bétoin, F. (279) 241
Feng, Y., see Tokuyama, S. (279) 93
Flower, R.J., see Harris, J.G. (279) 149
Franco-Cereceda, A., Barr, G., Öwall, A., Liska, J. and Lundberg, J.M.
Is endothelin-1 release at reperfusion of the ischaemic human heart due to cold-induced displacement of endothelin from binding sites? (279) 105
Fukao, M., see Hattori, Y. (279) 51

Goto, K., see Gyoda, Y. (279) 83
Guh, J.-H., Ko, F.-N., Yu, S.-M., Wu, Y.-C. and Teng, C.-M.
Pharmacological evaluation of *N*-methyl-actinodaphnine, a new vascular α-adrenoceptor antagonist, isolated from *Illigera luzonensis* (279) 33
Gyoda, Y., Tsukada, Y., Saito, A. and Goto, K.
Role of nitric oxide and neuropeptides in neurogenic vasodilatation of the guinea pig mesenteric artery (279) 83

Harris, J.G., Flower, R.J. and Perretti, M.
Alteration of neutrophil trafficking by a lipocortin 1 N-terminus peptide (279) 149
Hattori, Y., Kawasaki, H., Fukao, M. and Kanno, M.
Phorbol esters elicit Ca^{2+}-dependent delayed contractions in diabetic rat aorta (279) 51
Heyndrickx, J.-P., see Hodeige, D. (279) 25
Ho, I.K., see Tokuyama, S. (279) 93
Hodeige, D., Heyndrickx, J.-P., Chatelain, P. and Manning, A.
SR 33589, a new amiodarone-like antiarrhythmic agent: anti-adrenoceptor activity in anaesthetized and conscious dogs (279) 25
Horiai, H., see Terai, T. (279) 217
Horinouchi, T., see Koike, K. (279) 159

Igarashi, Y., see Van Ree, J.M. (279) 187
Imai, Y., see Schäfer, M. (279) 165

Jacobson, A.E., see Bowen, W.D. (279) R1

Kaku, S., Kawasaki, T., Sakai, Y., Taniuchi, Y., Yano, S., Suzuki, K.-i., Terazaki, C., Kawamura, K., Masuho, Y., Satoh, N., Takenaka, T., Yanagi, K. and Ohshima, N.
Antithrombotic effect of a humanized anti-GPIIb/IIIa monoclonal antibody, YM207, in a photochemically induced thrombosis model in monkeys (279) 115
Kanno, M., see Hattori, Y. (279) 51
Karcz-Kubicha, M. and Liljequist, S.
Evidence for an anxiogenic action of AMPA receptor antagonists in the plus-maze test (279) 171
Karhu, S., Perttula, S., Weckström, M., Kivistö, T. and Sellin, L.C.
Salicylaldoxime blocks K^+ and Ca^{2+} currents in rat cardiac myocytes (279) 7
Kawamura, K., see Kaku, S. (279) 115
Kawasaki, H., see Hattori, Y. (279) 51
Kawasaki, T., see Kaku, S. (279) 115
Keely, S.J., Stack, W.A., O'Donoghue, D.P. and Baird, A.W.
Regulation of ion transport by histamine in human colon (279) 203
Kita, Y., see Terai, T. (279) 217
Kivistö, T., see Karhu, S. (279) 7
Ko, F.-N., see Guh, J.-H. (279) 33
Koike, K., Horinouchi, T. and Takayanagi, I.
Possible mechanisms of β-adrenoceptor-mediated relaxation induced by noradrenaline in guinea pig taenia caecum (279) 159
Kuehne, M.E., see Bowen, W.D. (279) R1
Kumar, A., Raghubir, R. and Dhawan, B.N.
Possible involvement of nitric oxide in red nucleus stimulation-induced analgesia in the rat (279) 1
Kusunoki, T., see Terai, T. (279) 217

Lanz, I., see Amann, R. (279) 227
Lavarenne, J., see Bétoin, F. (279) 241
Lee, Y.-M., Sheu, J.-R. and Yen, M.-H.
BN-063, a newly synthesized adenosine A_1 receptor agonist, attenuates myocardial reperfusion injury in rats (279) 251
Lefebvre, R.A., see Briejer, M.R. (279) 123
Liljequist, S., see Karcz-Kubicha, M. (279) 171

Lin, W.-C.
Stimulatory effect of muscimol on gastric acid secretion stimulated by secretagogues in vagotomized rats under anesthesia (279) 43
Liska, J., see Franco-Cereceda, A. (279) 105
Lundberg, J.M., see Franco-Cereceda, A. (279) 105

Manning, A., see Hodeige, D. (279) 25
Marsden, C.A., see Wongwitdecha, N. (279) 99
Masuho, Y., see Kaku, S. (279) 115
Matsumoto, K., see Tanaka, K. (279) 143
Matsutani, S., see Tanaka, K. (279) 143
Meehan, S.M., see Schechter, M.D. (279) 59
Meldrum, B.S., see Millan, M.H. (279) 75
Millan, M.H., Chapman, A.G. and Meldrum, B.S.
Dual inhibitory action of enadoline (CI977) on release of amino acids in the rat hippocampus (279) 75
Miller, A.S. and Walker, J.M.
Effects of a cannabinoid on spontaneous and evoked neuronal activity in the substantia nigra pars reticulata (279) 179
Mironneau, C., see Pelet, C. (279) 15
Miura, T., see Yui, K. (279) 135

Nagase, M., see Yui, K. (279) 135
Neuilly, G., see Pelet, C. (279) 15

O'Donoghue, D.P., see Keely, S.J. (279) 203
Ohshima, N., see Kaku, S. (279) 115
Okuma, Y., see Yokotani, K. (279) 233
Ono, M., see Yui, K. (279) 135
Osumi, Y., see Yokotani, K. (279) 233
Öwall, A., see Franco-Cereceda, A. (279) 105

Paegelow, I., Werner, H. and Reißmann, S.
Effects of bradykinin and bradykinin analogues on spleen cells of mice (279) 211
Pelet, C., Mironneau, C., Rakotoarisoa, L. and Neuilly, G.
Angiotensin II receptor subtypes and contractile responses in portal vein smooth muscle (279) 15
Perretti, M., see Harris, J.G. (279) 149
Perttula, S., see Karhu, S. (279) 7

Raghubir, R., see Kumar, A. (279) 1
Rakotoarisoa, L., see Pelet, C. (279) 15
Reißmann, S., see Paegelow, I. (279) 211
Riekkinen, M., see Riekkinen Jr., P. (279) 65
Riekkinen Jr., P. and Riekkinen, M.
Effects of tetrahydroaminoacridine and nicotine in nucleus basalis and serotonin-lesioned rats (279) 65
Rust, C.J.J., see Van Ree, J.M. (279) 187

Saito, A., see Gyoda, Y. (279) 83
Sakai, Y., see Kaku, S. (279) 115
Satoh, N., see Kaku, S. (279) 115
Schäfer, M., Imai, Y., Uhl, G.R. and Stein, C.
Inflammation enhances peripheral μ-opioid receptor-mediated analgesia, but not μ-opioid receptor transcription in dorsal root ganglia (279) 165
Schechter, J.B., see Schechter, M.D. (279) 59
Schechter, M.D., Meehan, S.M. and Schechter, J.B.
Genetic selection for nicotine activity in mice correlates with conditioned place preference (279) 59
Schuligoi, R., see Amann, R. (279) 227
Schuurkes, J.A.J., see Briejer, M.R. (279) 123
Sellin, L.C., see Karhu, S. (279) 7
Sheu, J.-R., see Lee, Y.-M. (279) 251
Shibata, S., Tominaga, K. and Watanabe, S.
κ-Opioid receptor agonist protects against ischemic reduction of 2-deoxyglucose uptake in morphine-tolerant rats (279) 197
Shimazaki, T., see Terai, T. (279) 217
Shiokawa, Y., see Terai, T. (279) 217
Stack, W.A., see Keely, S.J. (279) 203
Stein, C., see Schäfer, M. (279) 165
Sugiyama, K., see Yui, K. (279) 135
Suzuki, K.-i., see Kaku, S. (279) 115

Takayanagi, I., see Koike, K. (279) 159
Takenaka, T., see Kaku, S. (279) 115
Tanaka, K., Matsutani, S., Matsumoto, K. and Yoshida, T.
Effect of thielocin A1β on bee venom phospholipase A_2-induced edema in mouse paw (279) 143
Taniuchi, Y., see Kaku, S. (279) 115
Teng, C.-M., see Guh, J.-H. (279) 33
Terai, T., Kita, Y., Kusunoki, T., Shimazaki, T., Ando, T., Horiai, H., Akahane, A., Shiokawa, Y. and Yoshida, K.
A novel non-xanthine adenosine A_1 receptor antagonist (279) 217
Terazaki, C., see Kaku, S. (279) 115
Tokuyama, S., Feng, Y., Wakabayashi, H. and Ho, I.K.
Ca^{2+} channel blocker, diltiazem, prevents physical dependence and the enhancement of protein kinase C activity by opioid infusion in rats (279) 93
Tominaga, K., see Shibata, S. (279) 197
Tsukada, Y., see Gyoda, Y. (279) 83

Uhl, G.R., see Schäfer, M. (279) 165

Vanderschuren, L., see Van Ree, J.M. (279) 187
Van Ree, J.M., Wolterink, G., Igarashi, Y., Vanderschuren, L., Wiegant, V.M., Rust, C.J.J. and Bruning, H.W.
Behavioral studies on the putative γ-type endorphin receptor using different antibodies (279) 187
Vilner, B.J., see Bowen, W.D. (279) R1

Wakabayashi, H., see Tokuyama, S. (279) 93
Walker, J.M., see Miller, A.S. (279) 179
Watanabe, S., see Shibata, S. (279) 197
Weckström, M., see Karhu, S. (279) 7
Werner, H., see Paegelow, I. (279) 211
Wiegant, V.M., see Van Ree, J.M. (279) 187
Wilcox, G., see Bétoin, F. (279) 241
Williams, W., see Bowen, W.D. (279) R1
Wolterink, G., see Van Ree, J.M. (279) 187
Wong, C.-L.
Effects of a histamine H_2 receptor agonist and antagonist on restraint-induced antinociception in female mice (279) 109
Wongwitdecha, N. and Marsden, C.A.
Isolation rearing prevents the reinforcing properties of amphetamine in a conditioned place preference paradigm (279) 99
Wu, Y.-C., see Guh, J.-H. (279) 33

Yanagi, K., see Kaku, S. (279) 115
Yano, S., see Kaku, S. (279) 115
Yen, M.-H., see Lee, Y.-M. (279) 251
Yokotani, K., Okuma, Y. and Osumi, Y.
Recombinant interleukin-1β inhibits gastric acid secretion by activation of central sympatho-adrenomedullary outflow in rats (279) 233
Yoshida, K., see Terai, T. (279) 217
Yoshida, T., see Tanaka, K. (279) 143
Yu, S.-M., see Guh, J.-H. (279) 33
Yui, K., Miura, T., Sugiyama, K., Ono, M. and Nagase, M.
Methamphetamine plus scopolamine potentiates behavioral sensitization and conditioning (279) 135